

# A note on coloring line arrangements

Eyal Ackerman\*

Rom Pinchasi†

## Abstract

We show that the lines of every arrangement of  $n$  lines in the plane can be colored with  $O(\sqrt{n/\log n})$  colors such that no face of the arrangement is monochromatic. This improves a bound of Bose et al. [1] by a  $\Theta(\sqrt{\log n})$  factor. Any further improvement on this bound will improve the best known lower bound on the following problem of Erdős: Estimate the maximum number of points in general position within a set of  $n$  points containing no four collinear points.

## 1 Introduction

Let  $\mathcal{A}$  be an arrangement of lines in the plane. Denote by  $\chi(\mathcal{A})$  the minimum number of colors required for coloring the lines of  $\mathcal{A}$  such that there is no monochromatic face in  $\mathcal{A}$ , that is, the boundary of every face is colored with at least two colors. Bose et al. [1] proved that  $\chi(\mathcal{A}) \leq O(\sqrt{n})$  for every *simple* arrangement  $\mathcal{A}$  of  $n$  lines, and that there are arrangements that require  $\Omega(\log n / \log \log n)$  colors. We improve their upper bound by a  $\Theta(\sqrt{\log n})$  factor, and extend it to not necessarily simple arrangements.

**Theorem 1.** *The lines of every arrangement of  $n$  lines in the plane can be colored with  $O(\sqrt{n/\log n})$  colors such that no face of the arrangement is monochromatic.*

A set of points in the plane is in *general position* if it does not contain three collinear points. Let  $\alpha(S)$  denote the maximum number of points in general position in a set  $S$  of points in the plane, and let  $\alpha_4(n)$  be the minimum of  $\alpha(S)$  taken over all sets  $S$  of  $n$  points in the plane with no four point on a line. Erdős pointed out that  $\alpha_4(n) \leq n/3$  and suggested the problem of determining or estimating  $\alpha_4(n)$ . Füredi [3] proved that  $\Omega(\sqrt{n \log n}) \leq \alpha_4(n) \leq o(n)$ .

We observe that any improvement of the bound in Theorem 1 would immediately imply a better lower bound for  $\alpha_4(n)$ . Indeed, suppose that  $\chi(\mathcal{A}) \leq k(n)$  for any arrangement of  $n$  lines, and let  $P$  be a set of  $n$  points, no four on a line. Let  $P^*$  be the dual arrangement of a slightly perturbed  $P$  (according to the usual point-line duality, see, e.g., [2, § 8.2]). Color  $P^*$  with  $k(n)$  colors such that no face is monochromatic, let  $S^* \subseteq P^*$  be the largest color class, and let  $S$  be its dual point set. Observe that the size of  $S$  is at least  $n/k(n)$  and it does not contain three collinear points, since the three lines that correspond to any three collinear points in  $P$  bound a face of size three in  $P^*$ .

## 2 Proof of Theorem 1

Let  $\mathcal{A}$  be an arrangement of  $n$  lines. We show that  $O(\sqrt{n/\log n})$  colors suffice for coloring the lines of  $\mathcal{A}$  such that no face in  $\mathcal{A}$  is monochromatic. Call a subset of lines *independent* if there is no face that is bounded just by lines from this subset. The proof is based on the following fact.

---

\*Department of Mathematics, Physics, and Computer Science, University of Haifa at Oranim, Tivon 36006, Israel. [ackerman@sci.haifa.ac.il](mailto:ackerman@sci.haifa.ac.il).

†Mathematics Department, Technion—Israel Institute of Technology, Haifa 32000, Israel. [room@math.technion.ac.il](mailto:room@math.technion.ac.il). Supported by BSF grant (grant No. 2008290).

**Theorem 2.** *There is an absolute constant  $c > 0$  such that every arrangement of  $n$  lines contains at least  $c\sqrt{n \log n}$  independent lines.*

We color the lines in  $\mathcal{A}$  such that no face is monochromatic by following the same method as in [1]. That is, we iteratively find a large subset of independent lines (whose existence is guaranteed by Theorem 2), color them with the same (new) color, and remove them from  $\mathcal{A}$ .

Clearly, this algorithm produces a valid coloring. We verify, by induction on  $n$ , that at most  $\frac{6}{c}\sqrt{n/\log n}$  colors are used in this coloring. We assume the bound is valid for all  $n \leq 256$  (by taking sufficiently small  $c > 0$ ). For  $n > 256$ , we have  $\log 4 < \frac{1}{4} \log n$ . Let  $i$  be the smallest integer such that after  $i$  iterations the number of remaining lines is at most  $n/4$ . Since in each of these iterations at least  $c\sqrt{\frac{n}{4} \log \frac{n}{4}} \geq c\sqrt{\frac{n}{8} \log n}$  vertices are removed,  $i \leq \frac{3n/4}{c\sqrt{\frac{n}{8} \log n}} \leq \frac{3}{\sqrt{2}c}\sqrt{n/\log n}$ . Therefore, by the induction hypothesis the number of colors that the algorithm uses is at most

$$i + \frac{6}{c}\sqrt{\frac{n}{4} \log \frac{n}{4}} < \frac{3}{\sqrt{2}c}\sqrt{\frac{n}{\log n}} + \frac{3}{c}\sqrt{\frac{n}{\log n - \frac{1}{4} \log n}} < \frac{3}{\sqrt{2}c}\sqrt{\frac{n}{\log n}} + \frac{2\sqrt{3}}{c}\sqrt{\frac{n}{\log n}} < \frac{6}{c}\sqrt{\frac{n}{\log n}} \quad \square$$

The proof of Theorem 2 is based on a result about independent sets in sparse hypergraphs. A *hypergraph*  $H = (V, E)$  consists of a set of vertices  $V$  and a set of edges  $E \subseteq 2^V$ . If the size of every edge is  $k$ , then  $H$  is  $k$ -uniform. For a set  $Z \subseteq V$  the *induced* hypergraph  $H[Z]$  consists of  $Z$  and the edges of  $H$  that are contained in  $Z$ . A set  $I \subseteq V$  is an *independent set* of  $H$ , if  $I$  contains no edge of  $H$ .

Kostochka et al. [4] proved that every  $n$ -vertex  $(k+1)$ -uniform hypergraph in which every  $k$  vertices are contained in at most  $d < n/(\log n)^{3k^2}$  edges contains an independent set of size  $\Omega\left(\left(\frac{n}{d} \log \frac{n}{d}\right)^{1/k}\right)$ . In fact, a careful look at their proof reveals the following result, that we stated for 3-uniform hypergraphs, since this is the case that we need.

**Theorem 3** ([4]). *Let  $H = (V, E)$  be an  $n$ -vertex 3-uniform hypergraph, such that every pair of vertices appear in at most  $d < n/(\log n)^{12}$  edges. Let  $X \subseteq V$  be a random set of vertices chosen independently with probability  $p = n^{-2/5}/(d \log \log \log n)^{3/5}$ , and let  $Z$  be an independent set chosen uniformly at random from the independent sets in  $H[X]$ . Then with high probability  $\mathbf{E}[|Z|] \geq \Omega(\sqrt{n \log n})$ .*

With Theorem 3 in hand we can now prove Theorem 2.

*Proof of Theorem 2:* Let  $L$  be a set of  $n$  lines and let  $\mathcal{A}$  be the corresponding arrangement. For a subset  $L' \subseteq L$  we say that a face  $f$  of  $\mathcal{A}$  is *bad* with respect to  $L'$  if all the lines bounding  $f$  are in  $L'$ .

Define a 3-uniform hypergraph  $H$  whose vertex set are the lines in  $\mathcal{A}$  and whose edge set consists of sets of three vertices such that the corresponding lines bound a face of size three in  $\mathcal{A}$ . Observe that every pair of lines can bound at most four faces of size three, therefore every pair of vertices in  $H$  appears in at most four edges. Let  $X$  be a random set of vertices (lines) chosen independently with probability  $p = n^{-2/5}/(4 \log \log \log n)^{3/5}$ . There are  $O(n^2)$  faces in  $\mathcal{A}$  and  $O(n)$  of them are of size two (since every line can bound at most four such faces). Therefore, the expected number of bad faces with respect to  $X$  of size different than three is  $O(p^2 n + p^4 n^2) = o(\sqrt{n \log n})$ .

It follows from Theorem 3 and Markov's inequality that with high probability the set of lines that correspond to  $X$  contains a subset  $Z$  of  $\Omega(\sqrt{n \log n})$  lines such that there are no faces of size three that are bad with respect to  $Z$ , and the number of bad faces of size different than three is  $o(\sqrt{n \log n})$ . By removing from  $Z$  a vertex from each bad face we obtain an independent set of lines of size  $\Omega(\sqrt{n \log n})$ .  $\square$

## References

- [1] P. Bose, J. Cardinal, S. Collette, F. Hurtado, S. Langerman, M. Korman, and P. Taslakian, Coloring and guarding arrangements, *28th European Workshop on Computational Geometry (EuroCG)*, March 19–21, 2012, Assisi, Perugia, Italy. Also available at: <http://arxiv.org/abs/1205.5162>
- [2] M. de Berg, O. Cheong, M. van Krefeld, M. Overmars, *Computational Geometry: Algorithms and Applications*, 3rd edition, Springer, 2008.
- [3] Z. Füredi, Maximal independent subsets in Steiner systems and in planar sets, *SIAM J. Disc. Math.* **4** (1991), 196–199.
- [4] A. Kostochka, D. Mubayi, J. Versatraete, On independent sets in hypergraphs, *Random Structures and Algorithms*, to appear.